



U.S. ENVIRONMENTAL PROTECTION AGENCY
 Office of Pesticide Programs
 Biopesticides and Pollution Prevention Division (7511P)
 1200 Pennsylvania Ave., N.W.
 Washington, D.C. 20460

EPA Reg. Number:

57538-64

Date of Issuance:

10/24/2017

NOTICE OF PESTICIDE:

Registration
 Reregistration
 (under FIFRA, as amended)

Term of Issuance:

Unconditional

Name of Pesticide Product:

Stimulate Plus Seed Germ

Name and Address of Registrant (include ZIP Code):

Stoller Enterprises, Inc.
 9090 Katy Freeway, Suite 400
 Houston, TX 77055

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Biopesticides and Pollution Prevention Division prior to use of the label in commerce. In any correspondence on this product, always refer to the above EPA Registration Number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA or the Act).

Registration is in no way to be construed as an endorsement or recommendation of this product by the U.S. Environmental Protection Agency (EPA). In order to protect health and the environment, the Administrator, on his or her motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under the Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is unconditionally registered in accordance with FIFRA section 3(c)(5) provided that you:

1. Submit and/or cite all data required for registration or registration review of your product when the EPA requires all registrants of similar products to submit such data.

Signature of Approving Official:

Andrew Bryceland, Team Leader
 Biochemical Pesticides Branch
 Biopesticides and Pollution Prevention Division (7511P)
 Office of Pesticide Programs

Date:

10/24/2017

2. Make the following labeling change before you release this product for shipment:
 - Revise the EPA Registration Number to read, “EPA Reg. No. 57538-64.”
3. Submit one (1) copy of the final printed labeling for the record before you release this product for shipment.

Should you wish to add/retain a reference to your company’s website on your label, then please be aware that the website becomes labeling under FIFRA and is subject to review by the EPA. If the website is false or misleading, the product will be considered to be misbranded and sale or distribution of the product is unlawful under FIFRA section 12(a)(1)(E). 40 CFR § 156.10(a)(5) lists examples of statements the EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the EPA find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA-approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance Assurance.

Your release for shipment of this product constitutes acceptance of these terms. If these terms are not complied with, this registration will be subject to cancellation in accordance with FIFRA section 6. A stamped copy of the labeling is enclosed for your records. Please also note that the record for this product currently contains the following acceptable Confidential Statement of Formula (CSF):

- Basic CSF dated 05/04/2017

Any CSFs other than those listed above are superseded.

If you have any questions, please contact Cody Kendrick of my team by phone at (703) 347-0468 or via email at kendrick.cody@epa.gov.

Sincerely,



Andrew Bryceland, Team Leader
Biochemical Pesticides Branch
Biopesticides and Pollution
Prevention Division (7511P)
Office of Pesticide Programs

Enclosure

STIMULATE™ PLUS SEED GERM

Sublabel A FOR AGRICULTURAL/COMMERCIAL USE

Sublabel B FOR HOMEOWNERS' GARDEN USE

ACTIVE INGREDIENTS:


Cytokinin (as kinetin)	0.05%
Gibberellic acid	0.16%
Indole-3-butyric acid.....	0.04%
Indole-3-acetic acid	0.04%

OTHER INGREDIENTS:	<u>99.71%</u>
Total	100.00%

This product contains 0.01538 grams cytokinin, 0.04920 grams gibberellic acid, 0.01230 grams indole-3-butyric acid, and 0.01230 indole-3-acetic acid per fluid ounce.

CONTAINS NON-PLANT FOOD INGREDIENTS:

- 0.05% Cytokinin
- 0.16% Gibberellic Acid
- 0.04% Indole-3-butyric Acid
- 0.04% Indole-3-acetic Acid



Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID	
If inhaled	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
If in eyes	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 seven days a week, 6:30 am to 4:30 pm Pacific Time (NPIC web site: www.npic.orst.edu).

See additional Precautionary Statements on left panel.

EPA Reg. No. 57538-

Read attached label before using.

EPA Est. No. 057538-TX-2,
57538-FL-1, 057538-IA-1

Not for Use in California

Manufactured and Guaranteed By
STOLLER ENTERPRISES, INC.
9090 Katy Freeway, Suite 400
Houston, Texas 77024 U.S.A.

Phone (713) 461-1493
Web: www.stollerusa.com E-Mail info@stollerusa.com

Density: 8.67 lbs/Gallon or 1.04 kg/Liter

- 1 Pint (1.08 lbs) 1 Quart (2.17) 1 Gallon (8.67 lbs)
- 2.5 Gallons (21.7 lbs) 5 Gallons (43.4 lbs) 55 Gallons (477 lbs)
- 4 Liters (4.16 kg) 10 Liters (10.4 kg) 20 Liters (20.8 kg) 208 Liters (216.32 kg)

STIMULATE™ PLUS SEED GERM

Sublabel A for AGRICULTURAL/COMMERCIAL USE

ACTIVE INGREDIENTS:

Cytokinin (as kinetin)	0.05%
Gibberellic acid	0.16%
Indole-3-butyric acid.....	0.04%
Indole-3-acetic acid	0.04%

OTHER INGREDIENTS:	<u>99.71%</u>
Total	100.00%

This product contains 0.01538 grams cytokinin, 0.04920 grams gibberellic acid, 0.01230 grams indole-3-butyric acid, and 0.01230 indole-3-acetic acid per fluid ounce.

CONTAINS NON-PLANT FOOD INGREDIENTS:

- 0.05% Cytokinin
- 0.16% Gibberellic Acid
- 0.04% Indole-3-butyric Acid
- 0.04% Indole-3-acetic Acid

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID	
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 seven days a week, 6:30 am to 4:30 pm Pacific Time (NPIC web site: www.npic.orst.edu).</p>	

See additional Precautionary Statements on left panel.

EPA Reg. No. 57538-

Read attached label before using.

EPA Est. Nos. 057538-TX-2,
57538-FL-1, 057538-IA-1

Not for Use in California

Manufactured and Guaranteed By
STOLLER ENTERPRISES, INC.
9090 Katy Freeway, Suite 400
Houston, Texas 77024 U.S.A.

Phone (713) 461-1493
 Web: www.stollerusa.com E-Mail info@stollerusa.com

Density: 8.67 lbs/Gallon or 1.04 kg/Liter

- 1 Pint (1.08 lbs) 1 Quart (2.17) 1 Gallon (8.67 lbs)
 2.5 Gallons (21.7 lbs) 5 Gallons (43.4 lbs) 55 Gallons (477 lbs)
 4 Liters (4.16 kg) 10 Liters (10.4 kg) 20 Liters (20.8 kg) 208 Liters (216.32 kg)

Sublabel A—Agricultural/Commercial Use
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if inhaled. Avoid breathing vapor or spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Remove and wash contaminated clothing before reuse. Wear the appropriate Personal Protective Equipment (PPE).

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Waterproof gloves are sufficiently chemical-resistant for this product. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear:

- long-sleeved shirt and long pants,
- chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride,
- shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS STATEMENT

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4.6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should: Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses, do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not apply when weather conditions favor drift from treated areas. Do not apply where runoff is likely to occur. Do not contaminate water when cleaning equipment or disposing of equipment washwater or rinsate. Exposed treated seed may be hazardous to birds and other wildlife. Treat only those seeds needed for immediate use and planting. Do not store excess treated seed beyond planting time. Dispose of all excess treated seed and seed packaging by burial away from streams and bodies of water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours unless wearing appropriate PPE.

Exception: If the product is soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, chemical-resistant gloves made of any waterproof material, and shoes plus socks.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter or allow others to enter treated areas until sprays have dried.

CHEMIGATION: APPLICATION AND CALIBRATION TECHNIQUES FOR SPRINKLER IRRIGATION

Apply this product only through the following types of irrigation systems: center pivot, traveler, big gun, motorized lateral move, end tow, side (wheel) roll, solid set, and hand move irrigation equipment. Do not apply through any other types of irrigation systems. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Experiment Station specialists, equipment manufacturers or other experts. Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person shall shut the system down and make necessary adjustments should the need arise.

A. Center Pivot, Traveler, Big Gun, Motorized Lateral Move, End Tow, and Side (Wheel) Roll Irrigation

Equipment: Operate system and injection equipment at normal pressures recommended by the manufacturer of injection equipment used. Fill tank of injection equipment with water. Operate system for one complete circle for center pivot or one complete run for the other recommended equipment, measuring time required, amount of water injected, and acreage contained in circle or run. Mix recommended amount of STIMULATE PLUS SEED GERM for acreage to be covered into same amount of water used during calibration and inject into system continuously for one revolution or run, but continue to operate irrigation system until STIMULATE PLUS SEED GERM has been cleared from last sprinkler head. Spray mixture in the chemical supply tank must be agitated at all times, otherwise settling and uneven application may occur.

B. Solid Set and Hand Move Irrigation Equipment: Determine acreage covered by sprinkler. Fill tank of injection equipment with water and adjust flow to use contents over a thirty to forty-five minute period. Mix desired amount of STIMULATE PLUS SEED GERM for acreage to be covered into quantity of water used during calibration and operate entire system at normal pressures recommended by the manufacturer of injection equipment used for amount of time established during calibration. Provide constant mechanical agitation in the mix tank to insure that STIMULATE PLUS SEED GERM will remain in suspension during the injection cycle. STIMULATE PLUS SEED GERM can be injected at the beginning or end of the irrigation cycle or as a separate application. Stop

injection equipment after treatment is completed and continue to operate irrigation system until STIMULATE PLUS SEED GERM is cleared from last sprinkler head.

SAFETY DEVICES

(1) The systems designated above must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow. (2) All pesticide injection pipelines must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. (3) The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. (4) The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. (5) The irrigation line or water pump must include a functional pressure switch, which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. (6) Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock. (7) Do not apply when wind speed favors drift beyond the area intended for treatment.

SYSTEMS CONNECTED TO PUBLIC WATER SOURCES

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops or, in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

For additional instructions on safety precautions, refer to statements (2), (3), (4) (6) and (7) in the section SAFETY DEVICES.

MODE OF ACTION

STIMULATE PLUS SEED GERM is a bio-stimulant containing plant growth regulators. Stimulate enhances plant growth and development by stimulating cell division, cell differentiation and enlargement, nutrient uptake and nutrient utilization. It is especially effective when applied with foliar fertilizer, but it is also compatible with pesticides.

MIXING INSTRUCTIONS: STIMULATE PLUS SEED GERM is water soluble and suitable for use in conventional liquid application systems including sprinkler irrigation systems. Shake Stimulate thoroughly and dilute in sufficient water to assure adequate, even coverage without producing excessive runoff. Agitate the spray mixture during application and apply within 12 hours of dilution. Stimulate can be applied tank mixed with most insecticides, fungicides, herbicides and foliar fertilizers but must be the last addition to the spray mixture.

APPLICATION INSTRUCTIONS

Apply STIMULATE PLUS SEED GERM (by ground or air) to foliage diluted in 2 to 10 gallons of water per acre. Larger volumes of water may be used if not associated with excessive runoff. For best results, apply in the early morning or late evening. When applying STIMULATE PLUS SEED GERM in a band or as a foliar-directed spray, reduce the application rate from the labeled broadcast rate in proportion to the percent of the field surface area covered by the foliar spray, but not below the minimum rate listed in the table.

VEGETABLES

Crop	No. of Ap- plications	Broadcast Spray		Band Spray		In Seed Furrow		Transplant Water		Spray Timing
		Oz/A	MI/H	Oz/A	MI/H	Oz/A	MI/H	Oz/A	MI/H	
Beans	1	0.8	56	0.4	28					Between the 3rd trifoliate leaf stage and flower bud formation.
	2-3	0.4	28	0.3	21	0.2-0.8	14-56			Begin at the 3rd trifoliate leaf stage and then at 7-10 day intervals.
	4-6	0.3	21	0.2	14					Begin at the 2nd trifoliate leaf stage and then at 7-14 day intervals.
Broccoli	3	0.6	42.5	0.4	28	0.2-0.8	14-56	0.2-0.8	14-56	Begin at the 4-5 leaf stage and then at 10-14 day intervals.
Brussel Sprouts	3	0.6	42.5	0.4	28	0.2-0.8	14-56	0.2-0.8	14-56	Begin at the 4-5 leaf stage and then at 10-14 day intervals.
Cabbage	3	0.6	42.5	0.4	28	0.2-0.8	14-56	0.2-0.8	14-56	Begin at the 4-5 leaf stage and then at 10-14 day intervals.
Cauliflower	3	0.6	42.5	0.4	28	0.2-0.8	14-56	0.2-0.8	14-56	Begin at the 4-5 leaf stage and then at 10-14 day intervals.
Corn, Sweet	1	0.8	56	0.5	35	0.2-0.8	14-56			2-6 leaf stage.
	2 or more	0.4	28	0.3	21					Begin at 2-6 leaf stage and then at 7-21 day intervals through the end of tasseling.
Cucumbers	1	0.8	56	0.4	28					Between flower bud initiation and first bloom.
	2-3	0.4	28	0.3	21	0.2-0.8	14-56	0.2-0.8	14-56	Begin at flower bud initiation and then at 7-10 day intervals.
	3-4	0.4	28	0.3	21					Begin at transplant, or at the 3-4 leaf stage for direct seeded, and then 7-10 day intervals.
Lettuce	3	0.6	42.5	0.4	28	0.2-0.8	14-56	0.2-0.8	14-56	Begin at the 4-5 leaf stage and then at 10-14 day intervals.
Melons	1	0.8	56	0.4	28					Between flower bud initiation and first bloom.
	2-3	0.4	28	0.3	21	0.2-0.8	14-56	0.2-0.8	14-56	Begin at flower bud initiation and then at 7-10 day intervals.
	4-6	0.3	21	0.2	14					Begin 2 weeks after emergence and then at 7-14 day intervals.
Onions	3	0.6	42.5	0.4	28	0.2-0.8	14-56			Begin 2 weeks after emergence and then at 10-14 day intervals.
Peppers	4-6	0.4	28	0.3	21	0.2-0.8	14-56	0.2-0.8	14-56	Begin at transplant, or at the 3-4 leaf stage for direct seeded, and then at 7-14 day intervals.
Potatoes	1	1.6	113.5	0.8	56	0.2-0.8	14-56			Tuber initiation.
	3	0.8	56	0.4	28					Begin at stolon formation (8-10 leaf stage) and then at 10-14 day intervals.
Squash	1	0.8	56	0.4	28	0.2-0.8	14-56			Between flower bud initiation and first bloom.
	2-3	0.4	28	0.3	21					Begin at flower bud initiation and then at 7-10 day intervals.
	4-6	0.3	21	0.2	14					Begin 2 weeks after emergence and then at 7-14 day intervals.
Tomatoes	1	0.8	56	0.4	28					Between flower bud initiation and first bloom.
	2-3	0.4	28	0.3	21	0.2-0.8	14-56	0.2-0.8	14-56	Begin at flower bud initiation and then at 7-10 day intervals.
	4-6	0.3	21	0.2	14					Begin 2 weeks after emergence, or at transplant, and then at 7-14 day intervals.

FIELD CROPS

Crop	No. of Ap- plications	Broadcast Spray		Band Spray		In Seed Furrow		Transplant Water		Spray Timing
		Oz/A	MI/H	Oz/A	MI/H	Oz/A	MI/H	Oz/A	MI/H	
Beets, sugar	1	1.6	113.5	0.8	56	0.2-0.8	14-56			6-8 leaf stage.
	2-3	0.8	56	0.4	28					Begin at the 2 leaf stage and then at 7-14 day intervals.
Canola	3	0.6	42.5	0.4	28	0.2-0.8	14-56			Begin at the 3-5 leaf stage and then at 10-14 day intervals.
Corn	1	0.8	56	0.5	35	0.2-0.8	14-56			2-6 leaf stage.
Cotton	1	0.8	56	0.4	28	0.2-0.8	14-56			Between flower initiation and final bloom.
	2-3	0.4	28	0.3	21					Early bloom and 7-14 days later.
	3-4	0.4	28	0.3	21					Begin at pinhead square and then at 7-10 day intervals.
Peanuts	4-6	0.4	28	0.3	21	0.2-0.8	14-56			Begin 30 days after planting and then at 7-14 day intervals.
Rice	1	0.8	56							2-5 leaf stage or panicle initiation.
	2	0.4	28							2-5 leaf stage or panicle initiation.
Sorghum	1	0.8	56	0.5	35	0.2-0.8	14-56			2-6 leaf stage.

Soybeans	1	0.8	56	0.5	35	0.2-0.8	14-56	3-7 trifoliolate leaf stage (V4-V8).
	2	0.4	28	0.3	21			3-7 trifoliolate leaf stage (V4-V8) and 10-17 days later.
Tobacco	2	0.8	56	0.5	35	0.2-0.8	14-56	1 st 40 days after planting 2 nd after topping
Wheat, Barley Oats, Rye	1-2	0.8	56	0.6	18	0.2-0.8	14-56	Start at tillering in the fall and/or spring and when 2 to 3 leaves form on main stem.

SMALL FRUITS, VINES AND TREE FRUITS

Bananas: Apply 0.4 to 0.8 fl. oz. /A (30 to 60 ml/hectare) in a band around the root mat and repeat every 10 to 14 days for a total of 10 applications.

Pineapple: Spray after transplanting with 0.4 to 0.8 fl. oz./100 gallons (12 to 24 ml/300 liters) and repeat every 10 to 14 days or apply in an irrigation system to provide 0.4 to 0.8 fl. oz./A (32.4 to 64.8 ml/hectare).

Strawberries: Begin sprays at first bloom at 0.4 fl oz/A (28.6 ml/hectare) broadcast or 0.2 fl oz/A (14.3 ml/hectare) as band sprays directed at the row. Repeat sprays at 2 to 4 week intervals for a total of 3 to 6 sprays.

Oranges: Apply STIMULATE PLUS SEED GERM at 0.1 to 0.2 pints/100 gallons (12.6 ml to 25.2 ml/100 liters) at first bloom and repeat at each flush of new growth.

YOUNG TREES AND ORNAMENTALS

Shrubs, Established: For increased vigor, rapid growth and healthy plant appearance, spray foliage with transplant solution (0.1 fl. oz. STIMULATE PLUS SEED GERM /2 gallons of water) to point of run-off two to three times per year.

Flowering Plants (Roses, Azaleas, etc.): 3-4 year old--using stock solution (0.1 fl. oz. STIMULATE PLUS SEED GERM /2 gallons of water) —Take 0.4 fl. oz. stock solution in 1 gallon of water and water in around root zone one time per year, preferably early spring.

Sod:* To improve growth, heavy rooting, broadcast 0.8 fl. oz./acre.

1. To speed up regrowth after harvest, broadcast 0.4 fl. oz./acre.
2. To boost with another 0.4 fl. oz./acre, broadcast six weeks later.

Turf:* For quick “tie down” after laying and to get turf off to a quick start, use STIMULATE PLUS SEED GERM as follows:

1. Broadcast 0.2 fl. oz./5,000 sq. ft. and water in.
2. Second application--30 days later—0.2 fl. oz./5,000 sq. ft. and water in.

*Do not apply this product through any type of irrigation system.

NOTICE: STIMULATE PLUS SEED GERM IS NOT A FERTILIZER. USE IN COMBINATION WITH A GOOD FERTILIZER PROGRAM WHERE INDICATED.

GOLF COURSES

Greens: Make an initial treatment with STIMULATE PLUS SEED GERM to promote root development and protect against “winter kill” using 0.2 fl. oz./5,000-7,000 sq. ft. Thereafter, use 0.1 fl. oz./green every 30 days.

Tees: Use STIMULATE PLUS SEED GERM at 0.05 fl. oz./1,200-1,500 sq. ft. of tee area every 30 days to maintain a healthy mass root growth.

Fairways: To establish the necessary root growth to fully utilize applied fertilizer, use STIMULATE PLUS SEED GERM at 0.8 fl. oz./acre two times the first year. Thereafter, use 0.8 fl. oz./acre one to two times a year.

TRANSPLANTS

STIMULATE PLUS SEED GERM MAY BE USED WHEN TRANSPLANTING ORNAMENTALS AND YOUNG TREES FOR A FAST AND HEALTHY START. Use 0.1 fl. oz./two gallons water stock solution as follows: At time of transplanting—

1. Bare (naked) roots - dip or spray with stock solution (0.1 fl. oz. STIMULATE PLUS SEED GERM /2 gallons of water).
2. Balled plants - spray ball at time of transplant.
3. Mist (not run-off) foliage lightly at time of transplant.
4. Using stock solution, apply 0.1 gallon STIMULATE PLUS SEED GERM in furrow/acre.

SEED TREATMENT

Use only on seeds for crops listed elsewhere on the label. Do not use treated seed for food, feed or oil purposes. If this product is intended for commercial seed treatment, treated seed must be labeled in accordance with the requirements of the Federal Seed Act and applicable State seed laws. An approved dye must be added to distinguish treated seed and prevent inadvertent use for food, feed, or oil purposes. If this product is intended for "at planting" use, treat only those seeds needed for immediate use and planting. Do not store excess treated seed beyond planting time. Dispose of excess treated seed by burial away from streams and bodies of water. A dye is not required.

For crops listed: Apply 0.1 to 0.4 fluid ounces per 100 lbs. of seed (.06 to 0.2 ml per kg). Dilute with water and mist spray on seed while mixing. Do not store seed wet as germination can be reduced if not planted soon after treatment.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Do not store in direct sunlight. Avoid freezing temperatures.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Handling: [Use label language appropriate for container size and type.]

Non-refillable containers. Do not reuse or refill this container. Clean container promptly after emptying.

Non-refillable container equal to or less than 5 gallons. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 120 seconds after the flow begins to drip. Fill the container $\frac{1}{4}$ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Non-refillable container greater than 5 gallons. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container $\frac{1}{4}$ full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure or accident, call CHEMTREC 1-800-424-9300

WARRANTY STATEMENT: To the extent consistent with applicable law, Stoller Enterprises, Inc. warrants that the product conforms to the description on the label and is reasonably fit for the purposes set forth on the label, when used according to directions under normal use conditions. Neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to the label instructions; the buyer assumes the risk of any such uses.

STIMULATE™ PLUS SEED GERM

Sublabel B FOR HOMEOWNERS' GARDEN USE

ACTIVE INGREDIENTS:

Cytokinin (as kinetin)	0.05%
Gibberellic acid	0.16%
Indole-3-butyric acid	0.04%
Indole-3-acetic acid	0.04%

OTHER INGREDIENTS:	<u>99.71%</u>
Total	100.00%

This product contains: 0.01538 grams cytokinin, 0.04920 grams gibberellic acid, 0.01230 grams indole-3-butyric acid, and 0.01230 indole-3-acetic acid per fluid ounce.

CONTAINS NON-PLANT FOOD INGREDIENTS:

- 0.05% Cytokinin
- 0.16% Gibberellic Acid
- 0.04% Indole-3-butyric Acid
- 0.04% Indole-3-acetic Acid

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.htm>

KEEP OUT OF REACH OF CHILDREN

CAUTION

FIRST AID	
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For emergency information concerning this product, call the National Pesticides Information Center (NPIC) at 1-800-858-7378 seven days a week, 6:30 am to 4:30 pm Pacific Time (NPIC web site: www.npic.orst.edu).</p>	

See additional Precautionary Statements on left panel.

EPA Reg. No. 57538-

Read attached label before using.

EPA Est. No. 057538-TX-2,
57538-FL-1, 057538-IA-1

Not for Use in California

Manufactured and Guaranteed By
STOLLER ENTERPRISES, INC.
9090 Katy Freeway, Suite 400
Houston, Texas 77024 U.S.A.

Phone (713) 461-1493
 Web: www.stollerusa.com E-Mail info@stollerusa.com

Density: 8.67 lbs/Gallon or 1.04 kg/Liter

- 1-Pint (1.08 lbs) 1 Quart (2.17) 1 Gallon (8.67 lbs)
 2.5 Gallons (21.7 lbs) 5 Gallons (43.4 lbs) 55 Gallons (477 lbs)
 4 Liters (4.16 kg) 10 Liters (10.4 kg) 20 Liters (20.8 kg) 208 Liters (216.32 kg)

Sublabel B For Homeowners' Garden Use
PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if inhaled. Avoid breathing vapor or spray mist. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Remove and wash contaminated clothing before reuse.

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm whether when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Do not enter or allow others to enter treated areas until sprays have dried.

MODE OF ACTION

STIMULATE PLUS SEED GERM is a bio-stimulant containing plant growth regulators. Stimulate enhances plant growth and development by stimulating cell division, cell differentiation and enlargement, nutrient uptake and nutrient utilization. It is especially effective when applied with foliar fertilizer, but it is also compatible with pesticides.

MIXING INSTRUCTIONS: STIMULATE PLUS SEED GERM is water soluble and suitable for use in conventional liquid application systems. Shake Stimulate thoroughly and dilute in sufficient water to assure adequate, even coverage without producing excessive runoff. Agitate the spray mixture during application and apply within 12 hours of dilution. If Stimulate is tank-mixed with other insecticides, fungicides, herbicides or foliar fertilizers, Stimulate must be the last addition to the spray mixture.

APPLICATION INSTRUCTIONS

Apply STIMULATE PLUS SEED GERM to foliage diluted in 1 gallon of water per 1000 sq. ft. Larger volumes of water may be used if not associated with excessive runoff. For best results, make early morning or late evening applications.

VEGETABLES

Crop	No. of Sprays	Amount/Gallon Per 1000 sq. ft.	Timing
Bean	3	0.2 teaspoons	Begin at the 3rd trifoliolate leaf and repeat every 7-10 days.
Broccoli, Brussels Sprout, Cauliflower	3	0.1 tablespoon	Begin at the 4 to 5 leaf-stage followed by 2 more applications at 10-14 day intervals.
Sweet Corn	2 or more	0.2 teaspoons	Begin at 2-6 leaf stage and then at 7-21 day intervals through the end of tasseling.
Cucumbers, Melon, Squash	3	0.2 teaspoons	Begin at flower bud initiation and then follow with 2 more sprays at 7-10 day intervals.
Lettuce	3	0.1 tablespoon	Begin at the 4-5 leaf stage and then 2 more sprays at 10-14 day intervals.

Onion	3	0.1 tablespoon	Begin 2 weeks after emergence and 2 more sprays at 10-14 day intervals.
Peppers	4	0.2 teaspoons	Begin at transplant or at the 3 to 4 leaf stage for direct seeded and repeat at 10-14 day intervals.
Potatoes	3	0.15 tablespoons	Begin at 8-10 leaf stage and then 2 more sprays at 7-10 day intervals.
Tomatoes	3	0.2 teaspoons	Begin at flower bud initiation and then 2 more sprays at 10-14 day intervals.

SMALL FRUITS, VINES AND TREE FRUITS

Crop	No. of Sprays	Amount/Gallon Per 1000 sq. ft.	Timing
Strawberries	3-6	0.2 teaspoons	Begin sprays at first bloom. Repeat at 2-4 week intervals for a total of 3-6 sprays.
Oranges	3-6	0.2 tablespoons	Spray to wet foliage at first bloom and repeat at each flush of new growth.
Shrubs, Established	2-3	0.4 tablespoons	Spray foliage to point of run-off 2-3 times a year.
Flower Plants (Roses, Azaleas, etc.)	1	0.1 teaspoon	In early spring, water in over root zone 1 time per year.

Turf*: To improve growth and heavy rooting, spray broadcast at 0.15 tablespoon per gallon per 1000 sq. ft. To speed up regrowth after cutting, spray broadcast 0.2 teaspoons per gallon of water per 1000 sq. ft.

Sod*: For quick “tie down” after laying and to get turf off to a quick start, use as follows:

1. Broadcast 0.2 fl. oz./5000 sq./ ft. and water in.
2. Make second application 30 days later and water in.

*Do not apply this product through any type of irrigation system.

Transplants: Prepare stock solution with 0.4 tablespoons of Stimulate Plus Seed Germ per 1 gallon of water. When transplanting with bare roots, dip in stock solution before planting. When transplanting with balled plants, spray ball at time of planting with stock solution. Mist foliage at time of transplant.

NOTICE: STIMULATE PLUS SEED GERM IS NOT A FERTILIZER. USE IN COMBINATION WITH A GOOD FERTILIZER PROGRAM WHERE INDICATED.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store unused product in original container only in a cool, dry area out of reach of children and animals. If container is damaged, place the container in a plastic bag. Do not store in direct sunlight. Avoid freezing temperatures.

Pesticide Disposal: If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

Container Handling: Non-refillable container. Do not reuse or refill this container. Place in trash or offer for recycling if available.

WARRANTY STATEMENT: To the extent consistent with applicable law, Stoller Enterprises, Inc. warrants that the product conforms to the description on the label and is reasonably fit for the purposes set forth on the label, when used according to directions under normal use conditions. Neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to the label instructions; the buyer assumes the risk of any such uses.